

Field Report

First record of a short-eared dog scavenging on an armadillo carcass

Holly O'Donnell^{1, 2}

¹ Fauna Forever, Avenida Aeropuerto, Km 1, La Joya, Puerto Maldonado, Madre de Dios, Peru.
Email: hollyodonnell146@gmail.com

² Wildlife Conservation Research Unit (WildCRU), Zoology, University of Oxford. Tubney House, Tubney OX13 5QL, UK

Keywords: Amazon; *Atelocynus microtis*; diet; Peru;

Abstract

The short-eared dog *Atelocynus microtis* is a solitary, medium sized canid endemic to South America. It is distributed across lowland Amazon rainforest habitat. Little is known of its ecology and there is limited information available on its diet, with records so far indicating a diet of fruit and live prey. Camera trap videos of a short-eared dog scavenging on the carcass of a nine-banded armadillo *Dasypus novemcinctus* over a period of four days were recorded along the Las Piedras river, Madre de Dios, Peru. This is the first record of scavenging by short-eared dogs.

Introduction

The short-eared dog *Atelocynus microtis* (Slater 1883) is a solitary, medium-sized South American canid of a monotypic genus (Leite-Pitman and Williams 2004, Koester et al. 2008, Leite-Pitman and Williams 2011). It is one of only two canid species known to exist in the Peruvian lowland Amazon rainforest, the other being the bush dog *Speothos venaticus* (Leite-Pitman and Williams 2004). They are the only two species of canids in the world which reside in dense rainforest habitat (Emmons 1997). Short-eared dogs are elusive and infrequently sighted (Leite-Pitman and Williams 2004, Leite-Pitman and Williams 2011), proving difficult to document (Pitman et al. 2003). Available data on the species remains scant (Sillero-Zubiri et al. 2004), with the species currently listed as Near Threatened (Leite-Pitman and Williams 2011).

The diet of the short-eared dog remains relatively undescribed (Leite-Pitman et al. 2008). Whilst skull morphometrics indicate that the species specialises in small vertebrates (Slater et al. 2009) it has also been described as omnivorous (Hunter and Caro 2008). Scat analysis showed short-eared dog diet to be comprised predominately of fish, but also included fruits, small mammals, frogs, and invertebrates (Pitman et al. 2003). Furthermore, food experiments with a semi-domesticated male determined food choice to be omnivorous but with a preference for meat (pers. comm. R. Leite-Pitman). Anecdotal, short-eared dogs have been noted to predate frogs and rodents such as spiny rats, agoutis *Dasypus sp.*, and pacas *Cuniculus sp.* (Emmons 1997). A camera trap recorded the first known instance of a short-eared dog carrying a caecilian (Apoda) in its mouth, which it was thought to have predated (Cisneros-Heredia and Mosquera 2010). The indigenous Sharanauas people of Alto Purus, Peru, recommended the use of peccary intestines as bait for the short-eared dog and it was successfully used to capture an individual in Los Amigos, Peru (pers. comm. R. Leite-Pitman). There are no published observations of the short-eared dog scavenging, and to the best of my knowledge this is the first record of this species scavenging.




Figure 1. Camera trap locations within the concession of the Las Piedras Amazon Center. Inset shows the location of the study area within the Madre de Dios region of south-eastern Peru.

The following is the established format for referencing this article:

O'Donnell, H. 2020. First record of a short-eared dog scavenging on an armadillo carcass. *Canid Biology & Conservation* 22(7): 25-28. URL: http://www.canids.org/CBC/22/Shorteared_dog_scavenging.pdf.

Results

Camera trap videos of a short-eared dog were recorded in June 2016 at the Las Piedras Amazon Center (LPAC) on the Lower Las Piedras River (12°04'13.1124"S, 69°29'38.1120"W) in the Madre de Dios region of south-eastern Peru (Figure 1). The habitat is lowland Amazon rainforest. The land is an ecotourism concession containing a field research station and conservation-tourism lodge. It covers an area of 44.60km² of low elevation flood-plain forest, higher elevation terra firme forest, bamboo forest patches, and palm swamp (aquajal). LPAC sits in a wider area of concessions, the vast majority of which are designated for the harvesting of Brazil nuts. There are several private conservation initiatives of similar size along the river within an approximately 20km radius. The closest state protected areas are the Tambopata National Reserve approximately 100 km to the south; the Madre de Dios Territorial Reserve approximately 100 km to the west (upriver from LPAC), and Manu National Park lying another 75 km beyond that. The Los Amigos private conservation concession (1,460km²) lies 85km south west of LPAC.

Figure 2



Figure 2. Still image from camera trap footage of a short-eared dog recorded at one of the permanent camera trap stations along a trail in the north-eastern area of the Las Piedras Amazon Center concession.

Two camera traps recorded six videos of the short-eared dog during a six-month survey focusing on a range of taxa. A Bushnell X-8 6MP camera trap was positioned on the intersection of two trails, located at 12°04'05.9880"S, 69°29'58.5960"W, from 3 March to 14 August 2016 (Figure 1). A lone short-eared dog was recorded three times during daylight hours, at 12:33, 12:33, and 12:34 on 21 June 2016 (video length 30 seconds; Figure 2). A second camera trap (Moultrie M1100i) was placed at the carcass of a nine-banded armadillo *Dasypus novemcinctus* at 12°04'06.4560"S, 69°30'06.1560"W on 12 June 2016 at 09:00 until 28 June 2016 at 09:11 (Figure 1). The carcass had no visible external wounds and appeared to have not yet been subject to scavenging. The trail close to which the carcass was found had been visited by the author 16 hours previously. The armadillo was therefore estimated to have likely died within the past 16 hours, or within a short time preceding that, due to the absence of the smell which led the author to detect the carcass. During the trapping period, three videos of a short-eared dog were recorded. The first video was recorded during daylight hours, likely on 21 June around 12:33h, however date and time settings were disrupted on the camera so exact time is unknown. In the first video the short-eared dog can be seen sniffing the area around the carcass for 29 seconds. Two further videos were recorded during the night three days later. In the first of the night-time videos the short-eared dog can be seen sniffing the carcass and moving its mouth on the carcass as though licking or tasting (Figure 3a, b). The final video shows the dog lifting a section of meat, which it drops again a few cm away (Figure 4a), with open mouth and teeth visible (Figure 4b). It is unclear whether the images show the same or two different individuals. It is only possible to determine sex of the individual in one video; it was determined to be a male. The short-eared dog was the only carnivore detected visiting or passing close to the carcass. Greater yellow-headed vultures *Cathartes melambrotus* and king vultures *Sarcoramphus papa* made up most of the images, nine-banded armadillo, collared peccary

Pecari tajacu, common opossum *Didelphis marsupialis*, and a green acouchi *Myoprocta microtis* were also recorded within the time-frame.

Discussion

Nine-banded armadillos are a wide-ranging, common prey species for carnivores in the Americas (Loughry et al. 2014). They are known to be the prey of several other carnivores in the Americas, all of which occur in the study area: jaguar *Panthera onca* (Núñez et al. 2000, Garla et al. 2001, Novack et al. 2005, Weckel et al. 2006, Harmsen et al. 2011); puma *Puma concolor* (Núñez et al. 2000); ocelot *Leopardus pardalis* (Wang 2002, Bianchi et al. 2010, de Oliveira et al. 2010); margay *Leopardus wiedii* (Wang 2002, Bianchi et al. 2011); and the bush dog (de Mello Beisiegel and Zuercher 2005, de Souza Lima et al. 2012). To-date there are no records to confirm that any armadillo species are the prey of short-eared dogs. At the time of the discovery of the armadillo carcass, the area was subject to a weather phenomenon known locally as a friaje, where cold weather fronts from Antarctica cause sudden wind, rain, and low temperatures in this area of the Amazon. Temperatures may drop by between 10 °C and 20 °C within a few hours, to lows of between 5 °C and 10 °C (Marengo, 1984). Cold weather may have initially slowed the rate of degradation of the carcass; however temperatures had risen to normal (~25 – 30°C) when the videos of the short-eared dog were recorded. The observed scavenging behaviour may be an opportunistic event as a direct result of the cold weather front, or it may be a more common occurrence. It is notable that the dog visits the carcass twice over four days, however further studies will be required to determine if this is characteristic dietary behaviour of the species because it is not possible to establish this from one record. Little is known of the ecology of the short-eared dog. Here I have presented what I believe to be the first recorded instance of this species scavenging an armadillo carcass. Due to the extreme difficulties associated with studying this species in the wild, opportunistic, and incidental observations can provide meaningful and useful ecological information. I call for greater publication of these observations to contribute to available information on this little-known species in order to inform conservation management decisions.

Figure 3a



Figure 3b



Figure 3. Still images from camera trap footage of a short-eared dog recorded at the armadillo carcass. The dog sniffs the carcass and moves its mouth on the carcass as though licking or tasting.



Figure 4. Still images from camera trap footage of a short-eared dog recorded at the armadillo carcass. (a) The dog's mouth is open, with teeth visible, and in contact with the carcass. (b) The dog lifts and moves a piece of armadillo with its teeth.

Acknowledgements

Thank you to Peru's Protected Area Authority (SERNANP), Forest and Wildlife Service (SERFOR), and Asociación para la Investigación y Desarrollo Integral (AIDER) for authorisations to undertake research at the field site. Thank you to Dr Chris Kirkby of Fauna Forever with whom I was employed as Mammal Research Coordinator at the time of the record. I am grateful to David Johnston, director of ARCAmazon, for hosting me at the Las Piedras Amazon Center. Thanks to Renata Leite-Pitman for her guidance with this manuscript; Rachel Rusen for donating one of the camera traps; Liselot Lange for the loan of a GPS; and Peter Coals and Cedric Tan for useful comments on an early draft.

References

- Bianchi, R.deC., Mendes, S.L. and de Marco Júnior, P. 2010. Food habits of the ocelot, *Leopardus pardalis*, in two areas in southeast Brazil. *Studies on Neotropical Fauna and Environment* 45(3): 111–119. DOI: [10.1080/01650521.2010.514791](https://doi.org/10.1080/01650521.2010.514791)
- Bianchi, R.deC., Rosa, A.F., Gatti, A. and Mendes, S.L. 2011. Diet of margay, *Leopardus wiedii*, and jaguarundi, *Puma yagouaroundi*, (Carnivora: Felidae) in Atlantic Rainforest, Brazil. *Zoologia (Curitiba)* 28(1): 127–132. DOI: [10.1590/S1984-46702011000100018](https://doi.org/10.1590/S1984-46702011000100018)
- Cisneros-Heredia, D.F. and Mosquera, D. 2010. First record of a canid (*Atelocynus microtis*) predating on a caecilian amphibian. *Reptilia* 24: 5–6.
- de Mello Beisiegel, B. and Zuercher, G.L. 2005. *Speothos venaticus*. *Mammalian Species* 783: 1–6. DOI: [10.1644/783.1](https://doi.org/10.1644/783.1)
- de Oliveira, T.G., Tortato, M.A., Silveira, L., Kasper, C.B., Mazim, F.D., Lucherini, M. and Sunquist, M.E. 2010. Ocelot ecology and its effect on the small-felid guild in the lowland neotropics. p. 559–580 in D.W. Macdonald and A.J. Loveridge (eds.), *Biology and conservation of wild felids*, Oxford University Press, Oxford, UK.
- de Souza Lima, E., Dematteo, K.E., Jorge, R.S.P., Jorge, M.L.S.P., Dalponte, J.C., Lima, H.S. and Klorfine, S.A. 2012. First telemetry study of bush dogs: Home range, activity and habitat selection. *Wildlife Research* 39(6): 512–519. DOI: [10.1071/WR11176](https://doi.org/10.1071/WR11176)
- Emmons, L. H. 1997. *Neotropical Rainforest Mammals: A Field Guide*. 2nd ed. The University of Chicago Press, Chicago, USA.
- Garla, R.C., Setz, E.Z.F., Gobbi, N., Dec, N., Garla, R.C., Paulista, U.E. and Paulo, S. 2001. Jaguar (*Panthera onca*) food habits in Atlantic rain forest of Southeastern Brazil. *Biotropica* 33(4): 691–696. DOI: [10.1111/j.1744-7429.2001.tb00226.x](https://doi.org/10.1111/j.1744-7429.2001.tb00226.x)
- Harmsen, B.J., Foster, R.J., Silver, S.C., Ostro, L.E.T. and Doncaster, C.P. 2011. Jaguar and puma activity patterns in relation to their main prey. *Mammalian Biology* 76(3): 320–324. DOI: [10.1016/j.mambio.2010.08.007](https://doi.org/10.1016/j.mambio.2010.08.007)
- Hunter, J. and Caro, T. 2008. Interspecific competition and predation in American carnivore families. *Ethology Ecology and Evolution* 20(4): 295–324. DOI: [10.1080/08927014.2008.9522514](https://doi.org/10.1080/08927014.2008.9522514)
- Koester, A.D., Azevedo, C.R., Vogliotti, A. and Duarte, J.M.B. 2008. Ocorrência de *Atelocynus microtis* (Sclater, 1882) na Floresta Nacional do Jamari, estado de Rondônia. *Biota Neotropica* 8(4): 231–234. DOI: [10.1590/S1676-06032008000400024](https://doi.org/10.1590/S1676-06032008000400024)
- Leite-Pitman, M.R.P. and Williams, R.S.R. 2004. The short-eared dog (*Atelocynus microtis*) p. 26–31 in C. Sillero-Zubiri, M. Hoffman and D.W. Macdonald (eds.) *Canids: foxes, wolves, jackals and dogs. Status Survey and Conservation Action Plan*. IUCN SSC Canid Specialist Group. Gland, Switzerland and Cambridge, UK.
- Leite-Pitman, R., Button, O., Hill, I., Lamy, K., & Nunnonca, E. (2008). Preferencia alimentaria del perro de orejas cortas (*Atelocynus microtis*). Presented at First Congress of Mastozoology in Peru, Universidad San Antonio de Abad del Cusco (UNSAAC), Cusco, Peru.
- Leite-Pitman, M.R.P. and Williams, R.S.R. 2011. *Atelocynus microtis*. The IUCN Red List of Threatened Species 2011. e.T6924A12814890. <https://www.iucnredlist.org/species/6924/12814890>. Downloaded 14 July 2018.
- Loughry, J., McDonough, C. and Abba, A.M. 2014. *Dasypus novemcinctus*. The IUCN Red List of Threatened Species 2014. e.T6290A47440785. <https://www.iucnredlist.org/species/6290/47440785>. Downloaded 14 July 2018.
- Marengo, J. 1984. Estudio Sinoptico-Climatico De Los Frijoles (Friegems) En La Amazonia Peruana. *Revista Forestal Del Perú* 12(1–2):1–26. Universidad Nacional Agraria La Molina, Peru. URL [http://cedinfor.lamolina.edu.pe/Articulos_RFP/Vol12_no1-2_84_\(16\)/vol12_art5.pdf](http://cedinfor.lamolina.edu.pe/Articulos_RFP/Vol12_no1-2_84_(16)/vol12_art5.pdf)
- Novack, A.J., Main, M.B., Sunquist, M.E. and Labisky, R.F. 2005. Foraging ecology of jaguar (*Panthera onca*) and puma (*Puma concolor*) in hunted and non-hunted sites within the Maya Biosphere Reserve, Guatemala. *Journal of Zoology* 267(2): 167–178. DOI: [10.1017/S0952836905007338](https://doi.org/10.1017/S0952836905007338)
- Núñez, R., Miller, B. and Lindzey, F. 2000. Food habits of jaguars and pumas in Jailsco, Mexico. *Journal of Zoology* 252(3): 373–379.
- Pitman, R., Beck, H. and Velazco, P. 2003. Mamíferos terrestres y arbóreos de la Selva Baja de la Amazonía Peruana entre los ríos Manu y Alto Purús. Pp 108–123 in R. Leite-Pitman, N. Pitman and P. Alvarez (eds.) *Alto Purús: Biodiversidad, conservación y manejo*, Center for Tropical Conservation, Duke University, USA.
- Sillero-Zubiri, C., Hoffman, M. and Macdonald, D.W. 2004. *Canids: foxes, wolves, jackals and dogs. Status survey and conservation action plan*. IUCN SSC Canid Specialist Group. Gland, Switzerland and Cambridge, UK.
- Slater, G.J., Dumont, E.R. and Van Valkenburgh, B. 2009. Implications of predatory specialization for cranial form and function in canids. *Journal of Zoology* 278(3): 181–188. DOI: [10.1111/j.1469-7998.2009.00567.x](https://doi.org/10.1111/j.1469-7998.2009.00567.x)

Wang, E. 2002. Diets of Ocelots (*Leopardus pardalis*), Margays (*L. wiedii*), and Oncillas (*L. tigrinus*) in the Atlantic rainforest in southeast Brazil. *Studies on Neotropical Fauna and Environment* 37: 207–212. DOI: [10.1076/snfe.37.3.207.8564](https://doi.org/10.1076/snfe.37.3.207.8564)

Weckel, M., Giuliano, W. and Silver, S. 2006. Cockscomb revisited : Jaguar diet in the Cockscomb Basin Wildlife Sanctuary. *Biotropica* 38(5): 687–690. DOI: [10.1111/j.1744-7429.2006.00190.x](https://doi.org/10.1111/j.1744-7429.2006.00190.x)

Biographical sketch

Holly O'Donnell works for the WildCRU, University of Oxford as a data assistant for the Trans-Kalahari Predator Project. She holds a PGDip in International Wildlife Conservation Practice from University of Oxford. Holly has a background working in the Peruvian Amazon, previously for the NGO Fauna Forever and now for Tamandua Expeditions.